



Canada's source for
HIV and hepatitis C
information

La source canadienne
de renseignements sur
le VIH et l'hépatite C

CATIE-News

CATIE's bite-sized HIV and hepatitis C news bulletins.

WHO guidelines encourage couples HIV testing and counselling and use of antiretroviral treatment for prevention

31 May 2012

In Canada, an estimated 83% of new HIV infections in 2008 occurred through sexual transmission, either among men who have sex with men (47%) or heterosexual men and women (36%). Among both groups a large number of these sexual HIV transmissions likely took place in the context of intimate relationships.

However, HIV prevention efforts in Canada focus mostly on changing the behaviour of an individual and often neglect the critical role partners may play in HIV transmission. Neglecting partners in HIV testing and counselling may be a missed opportunity for HIV prevention and treatment in Canada.

The World Health Organization (WHO) recently released "Guidance on Couples HIV Testing and Counselling," which outlines the important role of HIV testing and counselling for couples in HIV prevention, treatment and care.

The guidelines encourage countries to increase the offering of HIV testing and counselling to couples in addition to support for mutual disclosure of HIV status. In the guidelines, the WHO also recommends offering antiretroviral treatment to serodiscordant couples (in which one partner is HIV-positive and the other is HIV-negative) regardless of CD4 count in order to prevent HIV transmission.

Benefits of couples testing and counselling

The guidelines outline several potential benefits of offering HIV testing and counselling, along with support for mutual disclosure of HIV status, to couples. Many of these benefits are relevant to all couples, including those in which both partners are HIV-negative or both are HIV-positive.

Prevention of HIV transmission within couples and to external partners

In Canada, many people are unaware of their partner's, or their own, HIV status and therefore may unknowingly be at risk of becoming infected with HIV or transmitting HIV to their partner(s).

Research shows that when people become aware of their HIV infection, most take measures to reduce their risk of transmitting HIV to others. Furthermore, studies suggest couples who test and learn their status together are more likely to adopt preventive measures than those who test alone.

Couples HIV testing and counselling allows individuals to learn of their own status, and the status of their partner, and make decisions to prevent transmission together. In this context, counsellors can tailor prevention messages based on the couple's test results.

Increased uptake and adherence to antiretroviral therapy

A benefit of couples learning their test results together is that they can support each other, if one or both partners are HIV-positive, to access and adhere to ART for their own health and to prevent transmitting HIV to their partner or infant.

Also, the HIV-negative partner in a serodiscordant relationship may be able to protect themselves from HIV infection by taking antiretrovirals. This strategy is known as pre-exposure prophylaxis (PrEP).

Safer contraception and conception

Couples testing and counselling can help partners make informed decisions together with regards to reproductive health and family planning, including contraception and conception. Antiretroviral therapy or PrEP may be options to prevent HIV transmission when attempting to conceive.

Increased relationship cohesion and normalization of HIV status; decreased intimate partner violence and stigma

Testing and counselling for couples, with support for disclosure, may help couples come to terms with their partner's HIV status in a way that strengthens their relationship and can prevent stigma and violence. A counsellor can play an important role in creating a safe environment that can help ease tension and diffuse blame.

Recommendations on the use of antiretroviral treatment for HIV prevention

Over the past decade, evidence has been accumulating that shows antiretroviral treatment can reduce the risk that a person living with HIV transmits the virus to others. Last year, an important randomized controlled study known as HPTN 052 found that when HIV-positive partners in *heterosexual* serodiscordant relationships started ART early (with a CD4 count between 350 and 500 CD4 cells), HIV transmission was reduced by 96%.

Based on this study, and others, the WHO made two recommendations on the use of antiretroviral therapy to prevent HIV transmission between serodiscordant couples:

1. People with HIV in serodiscordant couples and who have started antiretroviral therapy (ART) for their own health should be advised that ART is also recommended to reduce HIV transmission to the uninfected partner.
2. HIV-positive partners with CD4 counts greater than 350 cells/ μ L in serodiscordant couples should be offered ART to reduce HIV transmission.
 1. People who are offered ART at this CD4 count should be told why they are being offered ART and that, once started, ART should be continued for life.
 2. ART should be offered to people with CD4 counts greater than 350 cells/ μ L for safer conception.

Currently, the WHO treatment guidelines do not recommend that people start ART for their own health if they have a CD4 count greater than 350 CD4 cells. However, in contrast to the WHO treatment guidelines, guidelines in high income countries generally recommend people living with HIV start treatment when their CD4 count falls below 500 cells. These recommendations are based on observational studies, which suggest that earlier treatment may improve the health of people living with HIV. Recently the US Department of Health and Human Services changed its guidelines to recommend treatment for all HIV-positive people who are ready to start regardless of their CD4 count.

The WHO guidelines on couples HIV testing and counselling emphasize that counsellors must ensure that HIV-positive individuals who are offered earlier treatment fully understand the purpose of this option and make an informed choice to start treatment. A number of factors must be taken into consideration when making the decision to start treatment earlier, such as the potential for side effects, the commitment to lifelong adherence, and the possibility of drug resistance developing due to poor adherence.

Counsellors must also ensure that couples are aware of other options for preventing HIV transmission, such as the consistent and correct use of male or female condoms. The following key messages with regards to condoms and antiretroviral therapy for prevention are outlined in the guidelines:

- It is possible for couples to stay HIV serodiscordant indefinitely if they consistently practice safer sex using condoms.
- Treatment for the HIV-positive partner is also highly effective in reducing the risk of transmission to the HIV-negative partner.
- Combined, treatment and consistent condom use are likely to offer greater protection than either one alone.

The WHO guidelines acknowledge that there is a lack of evidence with regards to how much ART reduces the risk of HIV transmission among populations other than heterosexual couples, such as gay couples and people who use injection drugs. However, the guidelines state that an "international scientific consensus is emerging that ART significantly reduces the risk of sexual HIV transmission of HIV regardless of the population or setting."

Pre-exposure prophylaxis to prevent HIV transmission among serodiscordant couples

In the past year several randomized controlled trials have found that HIV-negative individuals who take specific antiretroviral pills as pre-exposure prophylaxis are at reduced risk of HIV infection. The WHO is currently reviewing this evidence and plans to publish “rapid advice” on PrEP for heterosexual serodiscordant couples later this year. More detailed guidance will follow this advice.

Conclusion

Couples HIV testing and counselling provides an important opportunity to strengthen our HIV prevention efforts and should be offered in places in Canada where HIV testing is being done. Supporting linkages from HIV testing and counselling to appropriate treatment, care and support services will be important to fully realize the potential benefits of this couple-based intervention.

—James Wilton

Resources

[The Positive Side : Treatment benefits for all?](#)

[TreatmentUpdate: HPTN 052 - the trial that changed everything](#)

References

1. Guidance on couples HIV testing and counselling, including antiretroviral therapy for treatment and prevention in serodiscordant couples [Internet]. WHO. [cited 2012 May 22]. Available from: www.who.int/hiv/pub/guidelines/9789241501972/en/index.html
2. El-Bassel N, Gilbert L, Witte S et al. Couple-based HIV prevention in the United States: advantages, gaps, and future directions. *Journal of Acquired Immune Deficiency Syndromes*. 2010 Dec;55 Suppl2:S98-S101.
3. Marks G, Crepaz N, Senterfitt JW, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. *Journal of Acquired Immune Deficiency Syndromes*. 2005 Aug 1;39(4):446-53.

Produced By:



555 Richmond Street West, Suite 505, Box 1104
Toronto, Ontario M5V 3B1 Canada
Phone: 416.203.7122
Toll-free: 1.800.263.1638
Fax: 416.203.8284
www.catie.ca
Charitable registration number: 13225 8740 RR

Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to ensure they have the most current information. Users relying solely on this information do so entirely at their own risk. Neither CATIE nor any of its partners or funders, nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. Any opinions expressed herein or in any article or publication accessed or published or provided by CATIE may not reflect the policies or opinions of CATIE or any partners or funders.

Information on safer drug use is presented as a public health service to help people make healthier choices to reduce the spread of HIV, viral hepatitis and other infections. It is not intended to encourage or promote the use or possession of illegal drugs.

Permission to Reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by CATIE (the Canadian AIDS Treatment Information Exchange). For more information, contact CATIE at 1.800.263.1638.*

© CATIE

Production of this content has been made possible through a financial contribution from the Public Health Agency of Canada.

Available online at:

<http://www.catie.ca/en/catieneews/2012-05-29/who-guidelines-encourage-couples-hiv-testing-and-counselling-and-use-antiretrov>