



Canada's source for
HIV and hepatitis C
information

La source canadienne
de renseignements sur
le VIH et l'hépatite C

This fact sheet provides a snapshot of the HIV epidemic among people who inject drugs in Canada. It is one of a series of fact sheets on the epidemiology of HIV and hepatitis C.

All epidemiological information is approximate, based on the best available data. Most of the data in this fact sheet come from a population-specific surveillance system (I-Track), the *Summary: Estimates of HIV incidence, prevalence and Canada's Progress on Meeting the 90-90-90 HIV target, 2016*, and the *HIV/AIDS Epi Updates Chapter 1: Estimates of HIV prevalence and incidence in Canada, 2011* published by the Public Health Agency of Canada (PHAC). More information can be found in the section "[Where do these numbers come from?](#)" at the end of the fact sheet.

People who inject drugs represent 0.3% of the Canadian population.¹

The most recent population estimates for the number of people who inject drugs in Canada is from 2014.

According to 2014 national HIV estimates, there are 89,855 people who inject drugs in Canada. This represents 0.3% of the Canadian population aged 15 years and older.

People who inject drugs are 59 times more likely to get HIV than people who do not inject drugs.¹

The most recent estimates available that compare HIV incidence rates among key populations is from 2014. According to 2014 national estimates:

- The HIV incidence rate was 439 per 100,000 people who inject drugs.
- The HIV incidence rate was 7.5 per 100,000 people who do not inject drugs.
- People who inject drugs are 59 times more likely to get HIV than people who do not inject drugs.

Note: Because different methods were used to create the 2016 estimates, these 2014 estimates cannot be directly compared to the 2016 estimates.

14.3% of all new HIV infections in Canada may have been acquired through injection drug use (incidence).² **This varies considerably across Canada.**³

According to 2016 national HIV estimates:

- 14.3% of all new HIV infections in Canada may have been acquired through injection drug use. This means there were 310 new HIV infections in people who inject drugs in Canada. This includes:
 - 11.3% of new HIV infections where HIV status was attributed to injection drug use (244 new infections); and
 - 3% of new HIV infections where HIV status was attributed to the combined category of injection drug use or sex between men since both behaviours were reported at testing (66 new infections).

The most recent estimates available by region are for 2011. Based on estimates from 2011:

- The proportion of new HIV infections attributable to injection drug use varies across Canada:
 - 19% of new infections in British Columbia (72 new infections);
 - 17% in Alberta (42);
 - 79% in Saskatchewan (181);
 - 14% in Manitoba (16);
 - 9% in Ontario (120);
 - 11% in Quebec (80); and
 - 9% in the Atlantic provinces (3).

Note: Because different methods were used to create the 2016 estimates, these regional estimates from 2011 cannot be directly compared to the 2016 estimates.

A high proportion of new HIV infections (incidence) among women were likely due to injection drug use.²

According to 2016 national HIV estimates:

- 22% of the estimated new HIV infections in women were attributable to injection drug use. This compares to an estimated 11% of new infections among all Canadians attributable to injection drug use.

The number of new HIV infections (incidence) in people who inject drugs may have increased since 2014.²

According to 2016 national HIV estimates:

- The number of new HIV infections attributable to injection drug use in 2016 (244 new infections) is slightly higher than the number in 2014 (219 new infections).
- The number of new HIV infections attributable to the combined category of sex between men or injection drug use in 2016 (66 infections) is higher than in 2014 (47 new infections).

17.4% of people with HIV in Canada may have acquired their HIV infection through injection drug use (prevalence).² This varies considerably across Canada.³

According to 2016 national HIV estimates:

- 17.4% of people with HIV in Canada may have acquired their HIV infection through injection drug use (10,986 people). This includes:
 - 14.6% of people whose HIV status was attributed to injection drug use (9,204 people); and
 - 2.8% of people whose HIV status was attributed to the combined category of injection drug use or sex between men, since both behaviours were reported at testing (1,782 people).

The most recent estimates available by region are for 2011. Based on estimates from 2011:

- The proportion of people with HIV who may have acquired their HIV infection through injection drug use varies across Canada:
 - 34% in British Columbia (4,010 people)
 - 26% in Alberta (1,310 people)
 - 74% in Saskatchewan (1,550 people)
 - 19% in Manitoba (400 people)
 - 10% in Ontario (2,920 people)
 - 20% in Quebec (3,790 people)
 - 18% in the Atlantic provinces (180 people)

Note: Because different methods were used to create the 2014 estimates, these regional estimates from 2011 cannot be directly compared to the 2014 estimates.

Among people who use injection drugs, 11% have HIV and 68% either have or have had hepatitis C.⁴

According to I-Track (2010–2012):

- HIV prevalence among people who inject drugs in Canadian cities is approximately 11%.
- Approximately 68% of people who inject drugs had evidence of a current or past hepatitis C infection.
- Up to 10% of people who inject drugs may be co-infected with HIV and hepatitis C.

A high proportion of people who use drugs and report being HIV positive are engaged in care but fewer are currently on HIV treatment.³

According to I-Track (2010–2012):

- 95% of people who inject drugs and who self-report they are HIV positive are under the care of a doctor for their HIV.
- 66% of people who inject drugs and who self-report they are HIV positive are currently taking prescribed HIV drugs.

Key definitions

HIV prevalence—the number of people who are living with HIV at a point in time. Prevalence tells us how many people have HIV.

HIV incidence—the number of new HIV infections in a defined period of time (usually one year). Incidence tells us how many people are getting HIV.

Where do these numbers come from?

All epidemiological information is approximate, based on the best available data. Most of the data in this fact sheet come from a population-specific surveillance system (I-Track) or *HIV/AIDS Epi Updates Chapter 1: National HIV prevalence and incidence estimates for 2014* published by the Public Health Agency of Canada (PHAC).

Population-specific surveillance statistics

As part of the Federal Initiative to Address HIV/AIDS in Canada, PHAC monitors trends in HIV prevalence and associated risk behaviours among key vulnerable populations identified in Canada through population-specific surveillance systems. These surveillance systems, also known as the “Track” systems, are comprised of periodic cross-sectional surveys conducted at selected sites within Canada.

I-Track is the national surveillance system of people who inject drugs. For this surveillance system, information is collected directly from people who inject drugs through a questionnaire and a biological specimen sample collected for HIV and hepatitis C testing. The statistics provided in this fact sheet are for the years 2010 to 2012 from participating I-Track sites. Because the system only recruits voluntary participants from selected urban sites, the results do not represent all people who inject drugs across Canada.

Limitation—people from selected urban sites participated on a volunteer basis; therefore, the information presented does not represent all people who inject drugs in Canada.

National estimates of HIV prevalence and incidence

National HIV estimates are produced by PHAC and published every three years. Estimates of HIV prevalence and incidence are produced by PHAC using statistical methods which take into account some of the limitations of surveillance data (number of HIV diagnoses reported to PHAC), and also account for the number of people living with HIV who do not yet know they have it. Statistical modelling, using surveillance data and additional sources of information, allows PHAC to produce HIV estimates among those diagnosed and undiagnosed. The most recent estimates available are for 2014. The next set of estimates will be available in 2018 and will pertain to the year 2017.

Acknowledgements

We would like to thank the Surveillance and Risk Assessment Division of the Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada, for their helpful comments and expert review of CATIE epidemiology fact sheets.

References

1. [a.](#) [b.](#) Yang Q, Ogunnaike-Cooke S, Halverson J, et al. *Estimated national HIV incidence rates among key sub-populations in Canada, 2014*. Presented at *25th Annual Canadian Conference on HIV/AIDS Research (CAHR)*, 12–15 May 2016, Winnipeg, Canada. Abstract EPH3.5.
2. [a.](#) [b.](#) [c.](#) [d.](#) Public Health Agency of Canada. *Summary: Estimates of HIV Incidence, Prevalence and Canada’s Progress on Meeting the*

90-90-90 HIV targets, 2016. Public Health Agency of Canada, 2018. Available from: <https://www.canada.ca/en/public-health/services/publications/diseases-conditions/summary-estimates-hiv-incidence-prevalence-canadas-progress-90-90-90.html>.

3. [a. b. c.](#) Public Health Agency of Canada. *HIV/AIDS Epi Updates: National HIV Prevalence and Incidence Estimates for 2011*. Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada, 2014. Available from: <https://www.canada.ca/en/public-health/services/hiv-aids/publications/epi-updates/chapter-1-national-hiv-prevalence-incidence-estimates-2011.html>.
4. Public Health Agency of Canada. *Summary of Key Findings from I-Track Phase 3 (2010-2012)*. Surveillance and Epidemiology Division, Professional Guidelines and Public Health Practice Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada, 2012. Available from: <https://www.canada.ca/en/public-health/services/hiv-aids/publications/summary-key-findings-i-track-phase-3-2010-2012.html>.

Author(s): Challacombe L

Published: 2018

Produced By:



Canada's source for
HIV and hepatitis C
information

555 Richmond Street West, Suite 505, Box 1104
Toronto, Ontario M5V 3B1 Canada
Phone: 416.203.7122
Toll-free: 1.800.263.1638
Fax: 416.203.8284
www.catie.ca
Charitable registration number: 13225 8740 RR

Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to ensure they have the most current information. Users relying solely on this information do so entirely at their own risk. Neither CATIE nor any of its partners or funders, nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. Any opinions expressed herein or in any article or publication accessed or published or provided by CATIE may not reflect the policies or opinions of CATIE or any partners or funders.

Information on safer drug use is presented as a public health service to help people make healthier choices to reduce the spread of HIV, viral hepatitis and other infections. It is not intended to encourage or promote the use or possession of illegal drugs.

Permission to Reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by CATIE (the Canadian AIDS Treatment Information Exchange). For more information, contact CATIE at 1.800.263.1638.*

© CATIE

Production of this content has been made possible through a financial contribution from the Public Health Agency of Canada.

Available online at:
<http://www.catie.ca/en/fact-sheets/epidemiology/injection-drug-use-and-hiv-canada>