



Canada's source for
HIV and hepatitis C
information

La source canadienne
de renseignements sur
le VIH et l'hépatite C

From *HIV in Canada: A primer for service providers*

Sexual health promotion

Key Points

- Sexual health promotion is the delivery of strategies to promote sexual health.
- Promotion of sexual health behaviours can involve complex interventions that address issues related to empowerment, negotiation skills, self-esteem and sexual health knowledge.

The World Health Organization defines sexual health as:

“...a state of physical, emotional, mental and social well-being in relation to sexuality; it is not merely the absence of disease, dysfunction or infirmity. Sexual health requires a positive and respectful approach to sexuality and sexual relationships, as well as the possibility of having pleasurable and safe sexual experiences, free of coercion, discrimination and violence. For sexual health to be attained and maintained, the sexual rights of all persons must be respected, protected and fulfilled.” (WHO, 2006)

According to this definition, sexual health is an important part of everyone's overall health and well-being. It is essential that sexual health promotion activities encompass more than just prevention of HIV and other sexually transmitted infections but include activities that affect such things as healthy relationships, self-esteem, intimacy and pleasure.

Behavioural approaches to sexual health promotion attempt to motivate individuals and groups to change their behaviour through a range of educational, motivational, peer-group, skills-building and community normative approaches. Promotion of sexual health behaviours can involve complex interventions that address issues related to empowerment, negotiation skills, self-esteem and sexual health knowledge. Behavioural strategies to reduce the sexual transmission of HIV include attempts to delay the onset of first intercourse, decrease the number of sexual partners and increase the correct and consistent use of condoms.

Evidence suggests that the favourable behaviour changes seen in individuals during the first year following exposure to a behavioural prevention intervention can fade over time. At the population level, positive behaviour changes often fail to endure because these changes require a level of diligence — for example, consistent condom use — that is often difficult to maintain over the course of people's everyday lives and within their social contexts. This supports the need for ongoing messaging and support programming.

Resources

[Canadian guidelines for sexual health education](#) ^{PDF} – Public Health Agency of Canada

[SexandU.ca](#) – Society of obstetricians and gynecologists of Canada

Source

1. WHO. Defining sexual health: report of a technical consultation on sexual health, 28–31 January 2002. Geneva: World Health Organization; 2006. Available from: http://www.who.int/reproductivehealth/publications/sexual_health/defining_sexual_health.pdf ^{PDF}
2. World Health Organization. Sexual and reproductive health [Online]. Available from: http://www.who.int/reproductivehealth/topics/sexual_health/sh_definitions/en/index.html

Produced By:



Canada's source for
HIV and hepatitis C
information

555 Richmond Street West, Suite 505, Box 1104
Toronto, Ontario M5V 3B1 Canada
Phone: 416.203.7122
Toll-free: 1.800.263.1638
Fax: 416.203.8284
www.catie.ca
Charitable registration number: 13225 8740 RR

Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to ensure they have the most current information. Users relying solely on this information do so entirely at their own risk. Neither CATIE nor any of its partners or funders, nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. Any opinions expressed herein or in any article or publication accessed or published or provided by CATIE may not reflect the policies or opinions of CATIE or any partners or funders.

Information on safer drug use is presented as a public health service to help people make healthier choices to reduce the spread of HIV, viral hepatitis and other infections. It is not intended to encourage or promote the use or possession of illegal drugs.

Permission to Reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by CATIE (the Canadian AIDS Treatment Information Exchange). For more information, contact CATIE at 1.800.263.1638.*

© CATIE

Production of this content has been made possible through a financial contribution from the Public Health Agency of Canada.

Available online at:
<http://www.catie.ca/en/hiv-canada/4/4-2/4-2-3/4-2-3-1>