

From *Prevention in Focus*, Spring 2017



## Research Update: Youth who use drugs who cannot access addiction treatment are twice as likely to start injecting

By [Logan Broeckaert](#)

A Vancouver study<sup>1</sup> investigated the relationship between the accessibility of addiction treatment and the initiation of injection drug use among at-risk youth. This study found that youth who used drugs—cocaine, crack, heroin or methamphetamines— but who had never injected were twice as likely to start injecting after being unable to access addiction treatment.

The findings are from a cohort study of street-involved youth (aged 14–26) in Vancouver who had used drugs in the past 30 days. Participants had to have never injected drugs at study enrollment and to have at least one follow-up visit to be included in this study. Between September 2005 and May 2014, 462 participants met these criteria. Over the course of the study:

- 21% of participants disclosed that they had injected drugs for the first time (97 participants)
- 28% of participants disclosed that they had tried but were unable to access addiction treatment (129 participants)

Of the participants who were unable to access addiction treatment, 41% tried to access detox services, 35% treatment centres, 10% recovery houses, and 4% counsellors. Waiting lists were the most frequent barrier to treatment (66%); 18% of participants cited logistical barriers including location, hours and documentation requirements as barriers; and 6% for behavioural issues. Nine percent of participants were rejected by treatment programs for unspecified reasons.

The results of this study suggest that reducing barriers to treatment for youth who use drugs but who have never injected may reduce the likelihood that they will transition to injection drug use. This will avoid missed opportunities to prevent injection initiation and help to reduce drug-related harms like overdose and infection with HIV and hepatitis C.

### Reference

---

<sup>1</sup>. DeBeck K, Kerr T, Nolan S, et al. Inability to access addiction treatment predicts injection initiation among street-involved youth in a Canadian setting. *Substance Abuse Treatment, Prevention, and Policy* . 2016 Dec;11(1). Available from: <http://substanceabusepolicy.biomedcentral.com/articles/10.1186/s13011-015-0046-x>

### About the author(s)

**Logan Broeckaert** holds a Master's degree in History and is currently a researcher/writer at CATIE. Before joining

CATIE, Logan worked on provincial and national research and knowledge exchange projects for the Canadian AIDS Society and the Ontario Public Health Association.

## Produced By:



Canada's source for  
HIV and hepatitis C  
information

555 Richmond Street West, Suite 505, Box 1104  
Toronto, Ontario M5V 3B1 Canada  
Phone: 416.203.7122  
Toll-free: 1.800.263.1638  
Fax: 416.203.8284  
www.catie.ca  
Charitable registration number: 13225 8740 RR

## Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to ensure they have the most current information. Users relying solely on this information do so entirely at their own risk. Neither CATIE nor any of its partners or funders, nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. Any opinions expressed herein or in any article or publication accessed or published or provided by CATIE may not reflect the policies or opinions of CATIE or any partners or funders.

Information on safer drug use is presented as a public health service to help people make healthier choices to reduce the spread of HIV, viral hepatitis and other infections. It is not intended to encourage or promote the use or possession of illegal drugs.

## Permission to Reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by CATIE (the Canadian AIDS Treatment Information Exchange). For more information, contact CATIE at 1.800.263.1638.*

© CATIE

Production of this content has been made possible through a financial contribution from the Public Health Agency of Canada.

Available online at:  
<http://www.catie.ca/en/pif/spring-2017/research-update-youth-who-use-drugs-who-cannot-access-addiction-treatment-are-twice->