



Canada's source for
HIV and hepatitis C
information

La source canadienne
de renseignements sur
le VIH et l'hépatite C

From *Hepatitis C: An In-Depth Guide*

Nutrition on a budget

It is important that people living with hepatitis C eat a diet with adequate nutrients, including protein sources and vitamins and minerals from fruit and vegetables. This can be tough for people who have limited financial resources and often have to choose between food and shelter.

Key points of advice:

- Balance food intake with physical activity to maintain a healthy weight
- Avoid processed or fast food, which is high in fat and sugar

Some low-cost food choices:

Protein

Protein often comes from meat, which can be expensive. Choose alternate sources such as beans, lentils, peas or tofu. Canned tuna or salmon is also good and inexpensive (and low in fat). Protein can also be found in nuts, cheese, and eggs. If hunting or fishing is part of the culture or an activity done in the area, this can be a good way to get protein into your diet.

Carbohydrates

Pasta and rice are inexpensive and easy to prepare. When buying pasta, choose whole grain pasta or pasta high in fibre. Avoid white bread, donuts, cupcakes, cake and cookies. Look for whole grain breads and other baked goods on sale; buy “day old” bakery items and store them in the freezer until needed.

Fruits and vegetables

Canned or frozen fruits and vegetables are relatively inexpensive and store better than fresh. Different fruits and vegetables have different growing schedules—when a product is in season it is often cheaper and more widely available (for example, apples in the fall, berries in early summer), so take advantage of these deals. Following the natural growing schedule will also add variety to the diet.

Dairy

Powdered milk is less expensive than fresh milk and does not spoil. Orange juice with added calcium can be an alternative source of calcium for people who are lactose intolerant. Some vegetables like broccoli and kale also contain calcium.

Water

Try to drink lots of water every day. Water helps the body process nutrients and makes it easier for the liver to get rid of waste. Fruit juice is OK when it is hard to get clean drinking water. Alcohol, coffee and other drinks with caffeine (such as soda pop) don't count because they actually make the body lose water.

Community food programs

For people who run into trouble affording food, many communities have food bank programs. Different food banks have different policies about accessing their services. Other possible food access programs include drop-ins, community kitchens and meal programs.

Choosing a grocery store is important: convenience stores or small 24-hour grocery shops tend to be more expensive than discount grocery stores or supermarkets. There's a price attached to the convenience of 24-hour availability.

When possible, buying in bulk can save money. For example, breakfast cereals and oatmeal are much less expensive in bulk than in individual serving packets. This also applies to other staple items such as flour, sugar, dried fruits and nuts.

There are programs where local farmers provide people with fresh fruit and vegetables every two to four weeks for a small fee. These can be cheaper than going out to buy the vegetables individually.

Vitamins and supplements

Consulting a doctor before taking vitamin supplements will limit the potential for liver damage. A multivitamin-mineral supplement may be beneficial, particularly if the person eats irregularly or is not able to eat well. A physician can determine whether this is advisable.

No stove or fridge?

There are lots of foods that are healthy, keep for a while and don't require much cooking, like bread or bagels, peanut butter or nuts, granola bars, powdered milk, canned tuna or salmon, canned beans, raisins, bananas and apples. Try to keep some of these foods on hand.

Nutrition tips for severe liver damage (cirrhosis)

- Get advice from a doctor or dietitian (a healthcare professional who specializes in nutrition).
- Eat smaller meals every few hours. This may be easier to digest and will help keep energy levels up.
- Potentially limit intake of salt, if swelling of the stomach area (ascites) or legs (edema) is present.
- A multivitamin-mineral supplement is an option to discuss with a doctor. A version without iron may be needed.
- Try extra-strength liquid nutrition supplements, available from a pharmacy. Some examples are: Boost Plus Calories, Ensure Plus and Resource Plus.

Eating well when using drugs

Using drugs can suppress appetite or make it easy to forget to eat. Here are some things to do to get some nutrition while using and when taking a break.

- Drink high calorie meal replacement drinks, milkshakes, chocolate milk, fortified malted drinks or soymilk.
- Take a daily multivitamin-mineral supplement.
- Eat as well as possible when not high.
- Try to buy some groceries that last a long time. Stock up on peanut butter, oatmeal, powdered milk and canned stews and soups before spending money on drugs.

Detoxing or recovering from drug use?

Nutrition may be different while detoxing or recovering from drug use. Check out [Nutrition and recovery](#) for tips.

Finding a Dietitian

Registered dietitians have training in the science of food and nutrition. They can give specific advice about nutrition and Hep C. For more information on how to find a dietitian in your province or territory, see the [Dietitians of Canada Website](#).

Revised 2013.

Produced By:



Canada's source for
HIV and hepatitis C
information

555 Richmond Street West, Suite 505, Box 1104
Toronto, Ontario M5V 3B1 Canada
Phone: 416.203.7122
Toll-free: 1.800.263.1638
Fax: 416.203.8284
www.catie.ca
Charitable registration number: 13225 8740 RR

Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to ensure they have the most current information. Users relying solely on this information do so entirely at their own risk. Neither CATIE nor any of its partners or funders, nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. Any opinions expressed herein or in any article or publication accessed or published or provided by CATIE may not reflect the policies or opinions of CATIE or any partners or funders.

Information on safer drug use is presented as a public health service to help people make healthier choices to reduce the spread of HIV, viral hepatitis and other infections. It is not intended to encourage or promote the use or possession of illegal drugs.

Permission to Reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by CATIE (the Canadian AIDS Treatment Information Exchange). For more information, contact CATIE at 1.800.263.1638.*

© CATIE

Production of this content has been made possible through a financial contribution from the Public Health Agency of Canada.

Available online at:
<http://www.catie.ca/en/practical-guides/hepc-in-depth/living-hep-c/nutrition-budget>