



Canada's source for
HIV and hepatitis C
information

La source canadienne
de renseignements sur
le VIH et l'hépatite C

From *TreatmentUpdate* 227

Germany—experience with Maviret after licensure

Maviret is the brand name of a pill containing two anti-HCV drugs—glecaprevir and pibrentasvir. Maviret has been approved in Canada, the U.S. and the European Union for the treatment of hepatitis C virus (HCV) infection. For further information about how Maviret is used, see the [CATIE factsheet](#) on this drug.

Clinical trials tend to enroll relatively healthy people, so it is important to test newly licensed drugs in the people who attend clinics to get a better idea of the effects of the drug in a wider population.

Researchers in Germany collaborated in a large study with adults who had chronic HCV infection who either had not previously received HCV treatment or whose past treatment with interferon or the combination of sofosbuvir and ribavirin had failed. Data were collected between July 2017 and February 2018. In total, 321 people have completed treatment in this analysis and viral load information is still being collected. However, researchers confirmed that 96 people have been cured and 225 are currently being monitored.

The average profile of participants was as follows:

- 68% male, 32% female
- age - 47 years
- HCV viral load - 1.5 million IU/mL
- 90% had not previously been treated
- 71% had another co-existing health condition (such as cardiovascular disease)

The distribution of genotypes was as follows:

- GT 1a - 34%
- GT 1b - 17%
- GT 2 - 7%
- GT 3 - 35%
- GT 4 - 5%
- GT 5 or 6 - less than 1%

Most (98%) participants were treated with an eight-week regimen and the remaining 2% with a 12-week regimen.

Results

Three participants left the study prematurely, in two cases because of diarrhea or nausea. One participant stopped visiting the clinic. None of these people had serious symptoms of liver injury.

Among the remaining 93 patients who were cured and whose data were complete, common adverse events reported in the study were as follows:

- lack of energy - 9%
- headache - 8%

Such symptoms are commonly seen in people who have been treated with direct-acting antivirals (DAAs), regardless of the DAAs used, and resolve after a course of treatment has finished.

Severely abnormal blood test results were rare and occurred in three people. In two cases this was because of elevated liver enzymes, and in the third it was due to elevated levels of the waste product bilirubin. These abnormal blood test results were temporary.

The results from this observational study in Germany are very similar to the results seen with Maviret in clinical trials.

—Sean R. Hosein

REFERENCE:

Berg T, Stoehr A, Pangerl A, et al. First real-world data on safety and effectiveness of glecaprevir/pibrentasvir for the treatment of patients with chronic hepatitis C virus infection: data from the German Hepatitis C registry. *International Liver Congress*, 11-15 April 2018, Paris, France. Presentation GS-007

Produced By:



Canada's source for
HIV and hepatitis C
information

555 Richmond Street West, Suite 505, Box 1104
Toronto, Ontario M5V 3B1 Canada
Phone: 416.203.7122
Toll-free: 1.800.263.1638
Fax: 416.203.8284
www.catie.ca
Charitable registration number: 13225 8740 RR

Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to ensure they have the most current information. Users relying solely on this information do so entirely at their own risk. Neither CATIE nor any of its partners or funders, nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. Any opinions expressed herein or in any article or publication accessed or published or provided by CATIE may not reflect the policies or opinions of CATIE or any partners or funders.

Information on safer drug use is presented as a public health service to help people make healthier choices to reduce the spread of HIV, viral hepatitis and other infections. It is not intended to encourage or promote the use or possession of illegal drugs.

Permission to Reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by CATIE (the Canadian AIDS Treatment Information Exchange). For more information, contact CATIE at 1.800.263.1638.*

© CATIE

Production of this content has been made possible through a financial contribution from the Public Health Agency of Canada.

Available online at:

<http://www.catie.ca/en/treatmentupdate/treatmentupdate-227/hepatitis-c-virus/germany-experience-maviret-after-licensure>