



## MONOLAURIN (LAURIC ACID)

### What is it?

Monolaurin is a form of the fatty acid called lauric acid. It is found in human and coconut milk. In lab experiments, monolaurin has caused damage to a range of viruses, including HIV, herpes viruses and the bacteria that cause chlamydia and gonorrhea.

### What do HIV-positive people use this supplement for?

#### 1. To prevent and fight herpes outbreaks

Results from lab experiments suggest that monolaurin damages the herpes simplex virus, the virus that causes herpes sores. In those same experiments, monolaurin also attacked cytomegalovirus — a virus belonging to the herpes family commonly known as CMV — and to a lesser extent, HIV. These observations suggest that monolaurin may be most useful at suppressing outbreaks of herpes. To this end, most people take monolaurin in the form of capsules.

#### 2. To fight HIV

Some people obtain high levels of lauric acid by eating coconut milk or cream. Anecdotal reports from the United States suggest that taking the equivalent of three to four tablespoons of coconut oil daily significantly reduces viral load. On the other hand, anecdotal reports from Canada suggest that when taken daily for two years, coconut milk had no impact on CD4+ cell decline or the appearance of AIDS-related infections.

#### 3. To help speed recovery from bacterial infections such as chlamydia and gonorrhea.

Monolaurin has been shown to have anti-chlamydia and anti-gonorrhea effects in laboratory experiments involving cells. However, the effect of monolaurin in humans with these bacterial infections has not been documented in published studies.

### Available forms and usage

Monolaurin is supplied in 300 milligram capsules and is sold in health food stores. According to anecdotal reports, people who use monolaurin generally take between three and four capsules daily. Lauric acid can also be obtained by drinking coconut milk or by adding coconut milk or cream to liquid supplements, soups and other dishes.

### Cautions and concerns

The American Food and Drug Administration (FDA) recognizes monolaurin as safe enough to use as a food additive. Some people taking monolaurin have experienced drowsiness so users may prefer to take the capsules at bedtime. Finally, since we are unable to find reports of controlled studies using monolaurin, we can't be certain that monolaurin provides any clinical benefit.

### Credits

Author: Lori Lyons

Created: March 2000

Design: Renata Lipovitch



## References

Bergsson G, Arnfinnsson J, Karlsson SM et al. In vitro inactivation of Chlamydia trachomatis by fatty acids and monoglycerides. *Antimicrobial Agents and Chemotherapy* 1998;42(9):2290-2294.

Enig MG. Lauric oils as antimicrobial agents: theory of effect, scientific rationale, and dietary application as adjunct nutritional support for HIV-infected individuals. In: Watson RR ed., *Nutrients and Foods in AIDS*. New York: CRC Press, 1998.

Hamosh M, Peterson JA, Henderson TR, et al. Protective function of human milk: the milk fat globule. *Seminars in Perinatology* 1999;23(3):242-249.

Isaacs CE, Thormar H and Pessolano T. Membrane-disruptive effect of human milk: inactivation of enveloped viruses. *Journal of Infectious Diseases* 1986;154(6):966-971.

Isaacs CE, Kashyap S, Heird WC and Thormar H. Antiviral and antibacterial lipids in human milk and infant formula feeds. *Archives of Disease in Childhood* 1990;65(8):861-864.

Kristmundsdottir T, Arnadottir SG, Bergsson G and Thormar H. Development and evaluation of microbicidal hydrogels containing monoglyceride as the active ingredient. *Journal of Pharmaceutical Sciences* 1999;88(10):1011-1015.

Thormar H, Isaacs CE, Brown HR, et al. Inactivation of enveloped viruses and killing of cells by fatty acids and monoglycerides. *Antimicrobial Agents and Chemotherapy* 1987;31(1):27-31.

## Disclaimer

**Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV-related illness and the treatments in question.**

The Canadian AIDS Treatment Information Exchange (CATIE) in good faith provides information resources to help people living with HIV/AIDS who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

We do not guarantee the accuracy or completeness of any information accessed through or published or provided by CATIE. Users relying on this information do so entirely at their own risk. Neither CATIE nor Health Canada nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. The views expressed herein or in any article or publication accessed or published or provided by CATIE are solely those of the authors and do not reflect the policies or opinions of CATIE or the official policy of the Minister of Health Canada.

## Permission to reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by the Canadian AIDS Treatment Information Exchange (CATIE). For more information, contact CATIE at 1.800.263.1638.*

## Contact CATIE

**by telephone**

1.800.263.1638

416.203.7122

**by fax**

416.203.8284

**by e-mail**

info@catie.ca

**on the Web**

<http://www.catie.ca>

**by mail**

505-555 Richmond Street West

Box 1104

Toronto, Ontario

M5V 3B1

Canada



Funding has been provided by Health Canada, under the Canadian Strategy on HIV/AIDS.

