

# Lymphoma

## Summary

Lymphoma is a cancer that affects a part of the immune system called the lymphatic system. Like any cancer, the earlier lymphoma is diagnosed, the easier it is to treat. In general, HIV treatment (ART) significantly reduces the chance of people with HIV developing lymphoma. ART also strengthens the immune system and this can make the cancer easier to treat.

## About the lymphatic system

The lymphatic system is an extensive network of vessels that branches out into all parts of the body. The lymphatic system helps fight off infections and diseases. These vessels carry lymph, a clear fluid that contains cells of the immune system. Throughout the lymphatic system, small clusters of tissue, or lymph nodes, trap and filter germs. Larger groups of lymph nodes are found in the neck, armpits and groin. (These are the “swollen glands” you or your doctor may notice when you are ill.)

In addition to lymph nodes, other types of lymphatic tissue that are throughout your body include the tonsils, spleen, bone marrow and thymus, as well as patches of lymphatic tissue around your intestines. Because the immune system is so extensive, lymphoma can develop in, and spread to, almost any part of the body, including the spinal cord and brain.

## What is lymphoma?

Lymphoma is an umbrella term used to describe different cancers that affect the lymphatic system. There are two main types of lymphoma:

- Hodgkin’s lymphoma
- non-Hodgkin’s lymphoma

Lymphoma can be further divided into several subtypes based on the appearance of tumours when viewed under a microscope. These subtypes can include the following:

- diffuse large B-cell lymphoma (DLBCL)
- plasmablastic lymphoma
- primary effusion lymphoma

# FACT SHEET

**Published  
2016**

## CONTACT US

**by telephone**  
1-800-263-1638  
416-203-7122

**by fax**  
416-203-8284

**by e-mail**  
info@catie.ca

**by mail**  
555 Richmond Street West  
Suite 505, Box 1104  
Toronto ON M5V 3B1

- primary CNS lymphoma (this affects the brain and/or spinal cord)
- Burkitt lymphoma

## Who is at risk for lymphoma?

Large studies in the present era have found that HIV-positive people are at increased risk for these cancers. Researchers in North America, working with health-related information collected from nearly 90,000 HIV-positive and 200,000 HIV-negative people have estimated that by the age of 75 years about 5% of HIV-positive people will likely develop non-Hodgkin's lymphoma and about 1% will likely develop Hodgkin's lymphoma. The risk for these cancers is many times greater among HIV-positive people compared to people who are HIV negative.

It is very important to take ART every day, and achieve and maintain an undetectable viral load as this helps to strengthen the immune system. Although ART can make the amount of HIV in the blood undetectable and lengthen a person's survival, ART does not penetrate well into the lymphatic system. As a result, HIV can still infect cells of the immune system within lymph nodes and lymphatic tissues. These cells produce HIV-related proteins that can stimulate abnormal development and growth of cells of the immune system. Over time, one or more of these cells can turn into tumours. The cells that can turn into lymphoma tumours are mostly B-cells though, uncommonly, T-cells can also sometimes form lymphoma.

A relatively large study from Germany has investigated CD4+ and CD8+ cell counts a year prior to the diagnosis of lymphoma among HIV-positive people. Doctors found that in people with an undetectable viral load and who were taking ART everyday as directed, CD4+ cell counts decreased by an average of 168 cells 12 months before lymphoma was diagnosed. CD8+ cell counts decreased by about 352 cells 12 months before lymphoma was diagnosed. These decreasing cell counts may be a warning signal that lymphoma is developing. The figures we have listed were average changes in cell counts. In some patients the decreases in cell counts were relatively small. Cell counts may also decrease for other reasons and by

itself a cell count decrease does not mean that you have lymphoma.

Common members of the herpes virus family such as EBV (Epstein-Barr virus) have been linked to an increased risk for lymphoma.

## Symptoms

The most common symptom of lymphoma is swelling of the lymph nodes in the neck, armpits or groin. This swelling is usually painless. If you notice persistently swollen lymph glands, tell your doctor.

Other symptoms that often accompany swollen lymph nodes include:

- fever
- night sweats
- unintentional weight loss
- unexpected fatigue

Both swollen lymph nodes and the other symptoms mentioned above are common to many illnesses that can affect HIV-positive people. In other words, the presence of swollen lymph nodes with or without these symptoms does not always mean that you have cancer. However, it does mean that you need to see your doctor to have your health assessed.

If lymphoma develops in other parts of the body, it can cause symptoms related to those parts of the body. For example:

- Lymphoma in the gastrointestinal tract may cause abdominal pain or lead to an enlarged liver.
- Lymphoma in the mouth may cause pain and swelling in the mouth.
- Lymphoma in the brain can cause a headache, seizures or difficulty concentrating.

## Diagnosis

Lymphoma is diagnosed using a procedure called a biopsy. To do a biopsy, a doctor removes a small piece of tissue from the area of concern. The biopsied tissue is later examined under a

microscope to determine if lymphoma is present, what types of cells are involved, and how quickly the cancer cells are growing.

If a biopsy shows lymphoma, your doctor will likely recommend additional tests—such as an X-ray, or more sophisticated imaging techniques such as CT, MRI or PET scans.

These scans generally provide more information than an X-ray and help your doctor find out the size of the tumour(s), whether the cancer has spread and how far it may have spread. This process, called staging, will help determine the most suitable treatment.

In some cases an operation, called a laparotomy, may be done to allow your doctor to view internal organs and do more biopsies. A bone marrow biopsy and spinal tap may also be required to judge the extent of the lymphoma.

## Treatment

ART can make non-Hodgkin's lymphoma easier to treat. Since the 1990s, studies have found that HIV-positive people who take ART and who are later diagnosed with non-Hodgkin's lymphoma have been living longer than people who do not take ART. Your cancer doctor will develop a lymphoma treatment plan for you. The cancer treatment your doctor recommends will depend on the type of lymphoma you have, how far it has spread, how quickly it is growing, and your overall health—especially the health of your immune system. Your doctor will likely recommend one of the following approaches:

### A wait-and-see approach

If the lymphoma is small and growing slowly and has been diagnosed at a very early stage, your doctor may suggest a wait-and-see approach. This may seem surprising but bear in mind that treatment may not be initially necessary in some very early stages of cancer. Your doctor will continue to monitor your cancer closely and only advise therapy if and when the lymphoma starts to grow.

If your cancer is spreading more quickly, your doctor may recommend anti-cancer drugs (chemotherapy) or radiation therapy. When a tumour is developing and growing rapidly it is usually more susceptible to chemotherapy and radiation.

### Chemotherapy

Non-Hodgkin's lymphoma is usually treated with drugs (called chemotherapy or chemo) that kill cancer cells. The drugs may be taken as pills or injected into a vein (intravenous), into a muscle (intramuscular) or into the fluid that surrounds the spinal cord and brain. Side effects from chemotherapy are temporary and can include nausea, vomiting, fatigue, diarrhea, sores in the mouth, and hair loss. Your doctor may prescribe additional drugs to reduce the side effects. Your cancer nurse and pharmacist can help you find ways to cope with some of these temporary side effects.

### Radiation therapy

Radiation therapy may be used instead of, or in addition to, chemotherapy. This treatment uses radiation to help shrink tumours and destroy cancer cells.

Radiation therapy uses a machine outside the body to direct high-energy X-rays at a specific area of the body. The number of radiation treatments required varies from person to person, depending on the person's overall health and immune status, the location of the lymphoma, and how well the treatments are tolerated.

In a minority of cases, after it has been treated and resolved, lymphoma may recur. Clinical trials have found that bone marrow transplants can help HIV-positive people recover from lymphoma recurrence.

Unfortunately, both chemotherapy and radiation treatment temporarily weaken the immune system, and in some cases can put a person at risk of developing serious, sometimes life-threatening, infections called opportunistic infections. Your doctor may give you antibiotics and/or other drugs to reduce your risk of developing these infections.

ART has the potential to interact with chemotherapy, raising or lowering levels of chemotherapy drugs in your body. This can weaken its anti-cancer effect or cause you to experience enhanced side effects. Usually, a pharmacist at your cancer treatment centre will review your ART (and all the medicines that you are taking) to be sure that it does not interact with chemotherapy. Doctors who treat HIV-positive people who have cancer increasingly tend to ensure that their ART contains an integrase inhibitor as this group of drugs does not generally interact with chemotherapy.

## Outlook

Treatment can usually cause tumours to shrink and even disappear. As a result the signs and symptoms of the cancer can also disappear. Once high-resolution scans (CT, MRI or PET) cannot find any more tumours and your symptoms have resolved, ask your doctor if your cancer is in remission. People who have survived for five years in remission are generally considered to be cured. As ART has dramatically improved the overall health and survival of HIV-positive people, survival rates after lymphoma has gone into remission are very good. Note that, in a minority of cases, the lymphoma can return. If this happens, your doctor(s) can develop a new treatment plan and discuss options. Progress is being made in the field of cancer treatment and options for managing recurrence may be different from your initial treatment of cancer.

## References

- Dolcetti R, Ghoghini A, Caruso A, et al. A lymphomagenic role for HIV beyond immune suppression? *Blood*. 2016 Mar 17; 127(11):1403–9.
- Dolcetti R, Giagulli C, He W, et al. Role of HIV-1 matrix protein p17 variants in lymphoma pathogenesis. *Proceedings of the National Academy of Sciences USA*. 2015 Nov 17; 112(46):14331–6.
- Lorenzo-Redondo R, Fryer HR, Bedford T, et al. Persistent HIV-1 replication maintains the tissue reservoir during therapy. *Nature*. 2016 Feb 4;530(7588):51–6.
- Fletcher CV, Staskus K, Wietgreffe SW, et al. Persistent HIV-1 replication is associated with lower antiretroviral drug concentrations in lymphatic tissues. *Proceedings of the National Academy of Sciences USA*. 2014 Feb 11;111(6): 2307–12.

Uldrick TS, Little RF. How I treat classical Hodgkin lymphoma in patients infected with human immunodeficiency virus. *Blood*. 2015 Feb 19;125(8):1226–35.

Silverberg MJ, Lau B, Achenbach CJ, et al. Cumulative incidence of cancer among persons with HIV in North America: A cohort study. *Annals of Internal Medicine*. 2015 Oct 6; 163(7):507–18

Shiels MS, Pfeiffer RM, Besson C, et al. Trends in primary central nervous system lymphoma incidence and survival in the U.S. *British Journal of Haematology*. 2016 Aug;174(3):417–24.

Yanik EL, Achenbach CJ, Gopal S, et al. Changes in clinical context for Kaposi's sarcoma and non-Hodgkin lymphoma among people with HIV Infection in the United States. *Journal of Clinical Oncology*. 2016; *in press*.

Hoffmann C, Hentrich M, Gyllor D, et al. Hodgkin lymphoma is as common as non-Hodgkin lymphoma in HIV-positive patients with sustained viral suppression and limited immune deficiency: a prospective cohort study. *HIV Medicine*. 2015 Apr;16(4):261–4.

Hoffmann C, Schommers P, Wolf E, et al. CD4+ and CD8+ T-cell kinetics in aviremic HIV-infected patients developing Hodgkin or non-Hodgkin lymphoma. *AIDS*. 2016 Mar 13;30(5): 753-60.

Alvarnas JC, Le Rademacher J, Wang Y, et al. Autologous hematopoietic cell transplantation for HIV-related lymphoma: results of the BMT CTN 0803/AMC 071 trial. *Blood*. 2016 Aug 25;128(8):1050–8.

Barta SK, Samuel MS, Xue X, et al. Changes in the influence of lymphoma- and HIV-specific factors on outcomes in AIDS-related non-Hodgkin lymphoma. *Annals of Oncology*. 2015 May;26(5):958–66.

**Author(s):** Hosein SR, Koenig D, Maclean D, Sojé P

## Disclaimer

Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV- and hepatitis C-related illness and the treatments in question.

CATIE provides information resources to help people living with HIV and/or hepatitis C who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad a range of sources as possible. We strongly urge users to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

CATIE endeavours to provide the most up-to-date and accurate information at the time of publication. However, information changes and users are encouraged to consult as broad a range of sources as possible. Users relying on this information do so entirely at their own risk. Neither CATIE, nor any of its partners, funders, employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. The views expressed herein or in any article or publication accessed or published or provided by CATIE do not necessarily reflect the policies or opinions of CATIE nor the views of its partners and funders.

## Permission to reproduce

This document is copyrighted. It may be reprinted and distributed in its entirety for non-commercial purposes without prior permission, but permission must be obtained to edit its content. The following credit must appear on any reprint: *This information was provided by the Canadian AIDS Treatment Information Exchange (CATIE). For more information, contact CATIE at 1-800-263-1638.*

Production of this document has been made possible through a financial contribution from the Public Health Agency of Canada. The views expressed herein do not necessarily represent the views of the Public Health Agency of Canada.

CATIE fact sheets are available for free at [www.catie.ca](http://www.catie.ca)

## CONTACT US

### by telephone

1-800-263-1638  
416-203-7122

### by fax

416-203-8284

### by e-mail

[info@catie.ca](mailto:info@catie.ca)

### by mail

555 Richmond Street West  
Suite 505, Box 1104  
Toronto ON M5V 3B1

